

EAST: DERWENT EPO JPO USPAT 09/227,687 03/09/01

BRS	L5	309	424/93.1.icls.
BRS	L6	408	424/93.2.icls.
BRS	L7	336	424/93.21.icls.
BRS	L8	0	424/93.93.4.icls.
BRS	L9	160	424/93.4.icls.
BRS	L10	0	93.42.icls.
BRS	L12	288	424/234.1.icls.
BRS	L13	35	424/237.1.icls.
BRS	L11	4	424/93.42.icls.
BRS	L14	0	11 and gene
BRS	L15	1224	5 or 6 or 7 or 9 or 11 or 12 or 13
BRS	L16	909	15 and gene
BRS	L17	789	16 and recombinant
BRS	L18	406	15 and (gene with transform\$)
BRS	L19	147	18 and pathogen
BRS	L20	402	18 and (screen\$ or identif\$ or isolat\$)
BRS	L21	314	18 and ((screen\$ or identif\$ or isolat\$) with gene)
BRS	L22	2	5,981,182.pn.
BRS	L23	14	5801013.pn. 5871987.pn. 5885815.pn. 6174713.pn. 5656470.pn. 5798240.pn.
BRS	L24	12	tally-f\$.in.
BRS	L25	119	tao-j\$.in.
BRS	L26	32	wendler-p\$.in.
BRS	L27	11	connelly-g\$.in.
BRS	L28	78	gallant-d\$.in.
BRS	L29	0	24 and 25 and 26 and 27 and 28
BRS	L30	248	24 or 25 or 26 or 27 or 28
BRS	L31	6	30 and animal
BRS	L32	29	30 and (method with (determin\$ or isolat\$ or screen\$ or identif\$))
BRS	L33	0	30 and pro3
BRS	L34	0	30 and prors
BRS	L35	5	30 and aureus
BRS	L36	1308	514/44.icls.
BRS	L37	1112	36 and vivo
BRS	L38	469	37 and reporter
BRS	L39	120	38 and pathogen
BRS	L40	265	38 and ((transformed or recombinant) with cell)
BRS	L41	192	36 and ((ex adj2 vivo) and reporter)
BRS	L42	0	36 and ((ex adj2 vivo) with reporter)
BRS	L43	2	36 and ((ex adj2 vivo) with (lac or gfp or cat))
BRS	L44	122	36 and ((ex adj2 vivo) and reporter) and (gfp or lac or cat)
BRS	L45	17	44 and tet

+ Bajard-f.IN. + Tet

09/227,687  
Attach Paper # 15

(FILE 'HOME' ENTERED AT 18:27:20 ON 09 MAR 2001)

FILE 'MEDLINE, EMBASE, CAPLUS, BIOSIS' ENTERED AT 18:27:40 ON 09 MAR 2001

L1 485 S (TALLY, F?)/IN,AU  
L2 855 S (TAO, J?)/IN,AU  
L3 58 S (WENDLER, P?)/IN,AU  
L4 108 S (CONNELLY, G?)/IN,AU  
L5 72 S (GALLANT, C?)/IN,AU  
L6 492 S (GALLANT, D?)/IN,AU  
L7 0 S L1 AND L2 AND L3 AND L4 AND L6  
L8 1977 S L1 OR L2 OR L3 OR L4 OR L6  
L9 14 S L8 AND TET  
L10 6 DUPLICATE REMOVE L9 (8 DUPLICATES REMOVED)  
L11 15 S L8 AND (METHOD (S) (ISOLATION OR IDENTIFICATION OR SCREENING  
L12 11 DUPLICATE REMOVE L11 (4 DUPLICATES REMOVED)  
L13 0 S L8 AND PRO3  
L14 0 S L8 AND PC3844  
L15 0 S L8 AND PRORS  
L16 1 S L8 AND (INTRODUC? (S) (SUBJECT OR MAMMAL OR ANIMAL OR ORGANI  
L17 597 S ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?))  
AND  
L18 0 S L17 AND TET  
L19 126 S L17 AND INDUC?  
L20 75 S ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?))  
AND  
L21 0 S ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?))  
AND

Trying 3106016892...Open

Welcome to STN International! Enter x:x

LOGINID:sssptal636gx1

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International \* \* \* \* \*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Sep 29 The Philippines Inventory of Chemicals and Chemical  
Substances (PICCS) has been added to CHEMLIST  
NEWS 3 Oct 27 New Extraction Code PAX now available in Derwent  
Files  
NEWS 4 Oct 27 SET ABBREVIATIONS and SET PLURALS extended in  
Derwent World Patents Index files  
NEWS 5 Oct 27 Patent Assignee Code Dictionary now available  
in Derwent Patent Files  
NEWS 6 Oct 27 Plasdoc Key Serials Dictionary and Echoing added to  
Derwent Subscriber Files WPIDS and WPIX  
NEWS 7 Nov 29 Derwent announces further increase in updates for DWPI  
NEWS 8 Dec 5 French Multi-Disciplinary Database PASCAL Now on STN  
NEWS 9 Dec 5 Trademarks on STN - New DEMAS and EUMAS Files  
NEWS 10 Dec 15 2001 STN Pricing  
NEWS 11 Dec 17 Merged CEABA-VTB for chemical engineering and  
biotechnology  
NEWS 12 Dec 17 Corrosion Abstracts on STN  
NEWS 13 Dec 17 SYNTHLINE from Prous Science now available on STN  
NEWS 14 Dec 17 The CA Lexicon available in the CAPLUS and CA files  
NEWS 15 Jan 05 AIDSLINE is being removed from STN  
NEWS 16 Feb 06 Engineering Information Encompass files have new names  
NEWS 17 Feb 16 TOXLINE no longer being updated

NEWS EXPRESS FREE UPGRADE 5.0e FOR STN EXPRESS 5.0 WITH DISCOVER!  
(WINDOWS) NOW AVAILABLE

NEWS HOURS STN Operating Hours Plus Help Desk Availability

NEWS INTER General Internet Information

NEWS LOGIN Welcome Banner and News Items

NEWS PHONE Direct Dial and Telecommunication Network Access to STN

NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation  
of commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 18:27:20 ON 09 MAR 2001

=> FILE MEDLINE EMBASE CAPLUS BIOSIS

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.15

0.15

FILE 'MEDLINE' ENTERED AT 18:27:40 ON 09 MAR 2001

FILE 'EMBASE' ENTERED AT 18:27:40 ON 09 MAR 2001  
COPYRIGHT (C) 2001 Elsevier Science B.V. All rights reserved.

FILE 'CAPLUS' ENTERED AT 18:27:40 ON 09 MAR 2001  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 18:27:40 ON 09 MAR 2001  
COPYRIGHT (C) 2001 BIOSIS(R)

=> S (TALLY, F?)/IN,AU

'IN' IS NOT A VALID FIELD CODE  
'IN' IS NOT A VALID FIELD CODE  
L1 485 (TALLY, F?)/IN,AU

=> S (TAO, J?)/IN,AU

'IN' IS NOT A VALID FIELD CODE  
'IN' IS NOT A VALID FIELD CODE  
L2 855 (TAO, J?)/IN,AU

=> S (WENDLER, P?)/IN,AU

'IN' IS NOT A VALID FIELD CODE  
'IN' IS NOT A VALID FIELD CODE  
L3 58 (WENDLER, P?)/IN,AU

=> S (CONNELLY, G?)/IN,AU

'IN' IS NOT A VALID FIELD CODE  
'IN' IS NOT A VALID FIELD CODE  
L4 108 (CONNELLY, G?)/IN,AU

=> S (GALLANT, C?)/IN,AU

'IN' IS NOT A VALID FIELD CODE  
'IN' IS NOT A VALID FIELD CODE  
L5 72 (GALLANT, C?)/IN,AU

=> S (GALLANT, D?)/IN,AU

'IN' IS NOT A VALID FIELD CODE  
'IN' IS NOT A VALID FIELD CODE  
L6 492 (GALLANT, D?)/IN,AU

=> D HIS

(FILE 'HOME' ENTERED AT 18:27:20 ON 09 MAR 2001)

FILE 'MEDLINE, EMBASE, CAPLUS, BIOSIS' ENTERED AT 18:27:40 ON 09 MAR 2001

L1 485 S (TALLY, F?)/IN,AU  
L2 855 S (TAO, J?)/IN,AU  
L3 58 S (WENDLER, P?)/IN,AU  
L4 108 S (CONNELLY, G?)/IN,AU  
L5 72 S (GALLANT, C?)/IN,AU  
L6 492 S (GALLANT, D?)/IN,AU

=> S L1 AND L2 AND L3 AND L4 AND L6

L7 0 L1 AND L2 AND L3 AND L4 AND L6

=> S L1 OR L2 OR L3 OR L4 OR L6

L8 1977 L1 OR L2 OR L3 OR L4 OR L6

=> S L8 AND TET

L9 14 L8 AND TET

=> DUPLICATE REMOVE L9

DUPLICATE PREFERENCE IS 'MEDLINE, EMBASE, CAPLUS, BIOSIS'

KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):N

PROCESSING COMPLETED FOR L9

L10 6 DUPLICATE REMOVE L9 (8 DUPLICATES REMOVED)

=> D TI L10 1-6

L10 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2001 ACS

TI Tetracycline-inducible gene expression in gram-positive bacteria such as Staphylococcus and Bacillus

L10 ANSWER 2 OF 6 MEDLINE

DUPLICATE 1

TI Inhibition of protein synthesis occurring on tetracycline-resistant, TetM-protected ribosomes by a novel class of tetracyclines, the glycylyclines.

L10 ANSWER 3 OF 6 MEDLINE

DUPLICATE 2

TI Glycylyclines. 1. A new generation of potent antibacterial agents through modification of 9-aminotetracyclines.

L10 ANSWER 4 OF 6 MEDLINE

DUPLICATE 3

TI In vitro and in vivo antibacterial activities of the glycylyclines, a new class of semisynthetic tetracyclines.

L10 ANSWER 5 OF 6 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.

TI Characterization of pBFTM10, a clindamycin-erythromycin resistance transfer factor from Bacteroides fragilis.

L10 ANSWER 6 OF 6 CAPLUS COPYRIGHT 2001 ACS

TI Mechanisms of drug-resistance transfer in Bacteroides fragilis

=> D HIS

(FILE 'HOME' ENTERED AT 18:27:20 ON 09 MAR 2001)

FILE 'MEDLINE, EMBASE, CAPLUS, BIOSIS' ENTERED AT 18:27:40 ON 09 MAR 2001

L1 485 S (TALLY, F?)/IN,AU

L2 855 S (TAO, J?)/IN,AU

L3 58 S (WENDLER, P?)/IN,AU

L4 108 S (CONNELLY, G?)/IN,AU

L5 72 S (GALLANT, C?)/IN,AU

L6 492 S (GALLANT, D?)/IN,AU

L7 0 S L1 AND L2 AND L3 AND L4 AND L6

L8 1977 S L1 OR L2 OR L3 OR L4 OR L6

L9 14 S L8 AND TET

L10 6 DUPLICATE REMOVE L9 (8 DUPLICATES REMOVED)

=> S L8 AND (METHOD (S) (ISOLATION OR IDENTIFICATION OR SCREENING OR DETERMINING))

UNMATCHED LEFT PARENTHESIS 'AND (METHOD'

The number of right parentheses in a query must be equal to the number of left parentheses.

=> S L8 AND (METHOD (S) (ISOLATION OR IDENTIFICATION OR SCREENING OR DETERMINING))

L11 15 L8 AND (METHOD (S) (ISOLATION OR IDENTIFICATION OR SCREENING OR DETERMINING))

=> DUPLICATE REMOVE L11

DUPLICATE PREFERENCE IS 'MEDLINE, EMBASE, CAPLUS, BIOSIS'  
KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):N

PROCESSING COMPLETED FOR L11

L12 11 DUPLICATE REMOVE L11 (4 DUPLICATES REMOVED)

=> D TI L12 1-11

L12 ANSWER 1 OF 11 BIOSIS COPYRIGHT 2001 BIOSIS  
TI Enhancing drug discovery: Utilization of VITATM fluorescently labeled ligands in high throughput capillary electrophoresis screening.

L12 ANSWER 2 OF 11 MEDLINE DUPLICATE 1  
TI Detection of small-molecule enzyme inhibitors with peptides isolated from phage-displayed combinatorial peptide libraries.

L12 ANSWER 3 OF 11 BIOSIS COPYRIGHT 2001 BIOSIS  
TI Fungi for pitch reduction and their preparation.

L12 ANSWER 4 OF 11 BIOSIS COPYRIGHT 2001 BIOSIS  
TI Platform assay development strategy: Active-site directed peptides as tools for HTS.

L12 ANSWER 5 OF 11 BIOSIS COPYRIGHT 2001 BIOSIS  
TI Bacterial SecA as an antimicrobial target.

L12 ANSWER 6 OF 11 CAPLUS COPYRIGHT 2001 ACS  
TI Chemometric Labeling of Cereal Tissues in Multichannel Fluorescence Microscopy Images Using Discriminant Analysis

L12 ANSWER 7 OF 11 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V. DUPLICATE 2  
TI Membrane transport properties of mammalian oocytes: A micropipette perfusion technique.

L12 ANSWER 8 OF 11 CAPLUS COPYRIGHT 2001 ACS  
TI Analysis of petroleum acids in Dushanzi distillate

L12 ANSWER 9 OF 11 CAPLUS COPYRIGHT 2001 ACS  
TI T4 radioimmunoassay of dried blood samples on filter paper and its clinical application

L12 ANSWER 10 OF 11 MEDLINE DUPLICATE 3  
TI Differentiation of Bacteroides ovatus and Bacteroides thetaiotaomicron by means of bacteriophage.

L12 ANSWER 11 OF 11 BIOSIS COPYRIGHT 2001 BIOSIS  
TI QUANTITATIVE ISOLATION OF RADIO LABELED METABOLITES WITHOUT CHROMATOGRAPHY  
MEASUREMENTS OF THE BIOSYNTHESIS OF PURINES PYRIMIDINES AND UREA IN ISOLATED HEPATOCYTES.

L12 ANSWER 1 OF 11 BIOSIS COPYRIGHT 2001 BIOSIS  
ACCESSION NUMBER: 2000:540761 BIOSIS  
DOCUMENT NUMBER: PREV200000540761  
TITLE: Enhancing drug discovery: Utilization of VITATM  
fluorescently labeled ligands in high throughput capillary  
electrophoresis screening.  
AUTHOR(S): Finn, J. (1); Glicksman, M. (1); Riera, T. (1); Gallant,  
P.  
(1); **Tao, J. (1)**; Chapple, J. (1); Dunayevskiy,  
Y.; Hughes, D.  
CORPORATE SOURCE: (1) Cubist Pharmaceuticals, Inc., Cambridge, MA USA  
SOURCE: Abstracts of the Interscience Conference on Antimicrobial  
Agents and Chemotherapy, (2000) Vol. 40, pp. 226. print.  
Meeting Info.: 40th Interscience Conference on  
Antimicrobial Agents and Chemotherapy Toronto, Ontario,  
Canada September 17-20, 2000  
DOCUMENT TYPE: Conference  
LANGUAGE: English  
SUMMARY LANGUAGE: English

L12 ANSWER 2 OF 11 MEDLINE DUPLICATE 1  
ACCESSION NUMBER: 2000130935 MEDLINE  
DOCUMENT NUMBER: 20130935  
TITLE: Detection of small-molecule enzyme inhibitors with  
peptides  
isolated from phage-displayed combinatorial peptide  
libraries.  
AUTHOR: Hyde-DeRuyscher R; Paige L A; Christensen D J;  
Hyde-DeRuyscher N; Lim A; Fredericks Z L; Kranz J; Gallant  
P; Zhang J; Rocklage S M; Fowlkes D M; **Wendler P A**  
; Hamilton P T  
CORPORATE SOURCE: Novalon Pharmaceutical Corporation, Durham, NC 27703,  
USA.  
SOURCE: CHEMISTRY AND BIOLOGY, (2000 Jan) 7 (1) 17-25.  
Journal code: CNA. ISSN: 1074-5521.  
PUB. COUNTRY: ENGLAND: United Kingdom  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 200005  
ENTRY WEEK: 20000503

AB BACKGROUND: The rapidly expanding list of pharmacologically important  
targets has highlighted the need for ways to discover new inhibitors that  
are independent of functional assays. We have utilized peptides to detect  
inhibitors of protein function. We hypothesized that most peptide ligands  
identified by phage display would bind to regions of biological  
interaction in target proteins and that these peptides could be used as  
sensitive probes for detecting low molecular weight inhibitors that bind  
to these sites. RESULTS: We selected a broad range of enzymes as targets  
for phage display and isolated a series of peptides that bound  
specifically to each target. Peptide ligands for each target contained  
similar amino acid sequences and competition analysis indicated that they  
bound one or two sites per target. Of 17 peptides tested, 13 were found

to  
be specific inhibitors of enzyme function. Finally, we used two peptides  
specific for Haemophilus influenzae tyrosyl-tRNA synthetase to show that  
a

simple binding assay can be used to detect small-molecule inhibitors with  
potencies in the micromolar to nanomolar range. CONCLUSIONS: Peptidic  
surrogate ligands identified using phage display are preferentially  
targeted to a limited number of sites that inhibit enzyme function. These  
peptides can be utilized in a binding assay as a rapid and sensitive  
**method** to detect small-molecule inhibitors of target protein

function. The binding assay can be used with a variety of detection systems and is readily adaptable to automation, making this platform ideal for high-throughput **screening** of compound libraries for drug discovery.

L12 ANSWER 3 OF 11 BIOSIS COPYRIGHT 2001 BIOSIS  
ACCESSION NUMBER: 2000:279564 BIOSIS  
DOCUMENT NUMBER: PREV200000279564  
TITLE: Fungi for pitch reduction and their preparation.  
AUTHOR(S): Farrell, Roberta L. (1); Hadar, Yitzhak; **Wendler, Philip A.**; Zimmerman, Wendy  
CORPORATE SOURCE: (1) Watertown, MA USA  
ASSIGNEE: Clariant Finance (BVI) Limited, Tortola, British Virgin Islands  
PATENT INFORMATION: US 5998197 December 07, 1999  
SOURCE: Official Gazette of the United States Patent and Trademark Office Patents, (Dec. 7, 1999) Vol. 1229, No. 1, pp. No pagination. e-file..  
ISSN: 0098-1133.  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
AB Ascospores of wood-penetrating, pitch-grading fungi of the class of Ascomycotina and Deuteromycotina, eg. Ophiostromas, may be screened to provide fungi combining the properties of good growth on non-sterile wood substrates and minimized or even enhanced brightness effects for use in pitch reduction of wood substrates, eg. logs and wood chips. A new and improved method of isolating such ascospores involving effective suspension in an oil consumable by the fungus, eg. a vegetable oil, and then treatment of the oil with a dispersing agent is also disclosed.

L12 ANSWER 4 OF 11 BIOSIS COPYRIGHT 2001 BIOSIS  
ACCESSION NUMBER: 1999:258890 BIOSIS  
DOCUMENT NUMBER: PREV199900258890  
TITLE: Platform assay development strategy: Active-site directed peptides as tools for HTS.  
AUTHOR(S): **Wendler, P. (1)**; Gallant, P. (1); Kranz, J. (1); Lim, A. (1); Namchuk, M. (1); Zhang, J. (1); Rocklage, S. (1); Deruyscher; Paige, L.; Hyde-Deruyscher, N.; Hamilton, P.; Fredericks, Z.  
CORPORATE SOURCE: (1) Cubist Pharmaceuticals, Inc., Cambridge, MA USA  
SOURCE: Abstracts of the Interscience Conference on Antimicrobial Agents and Chemotherapy, (1998) Vol. 38, pp. 274.  
Meeting Info.: 38th Interscience Conference on Antimicrobial Agents and Chemotherapy San Diego, California, USA September 24-27, 1998 American Society for Microbiology  
DOCUMENT TYPE: Conference  
LANGUAGE: English

=> D HIS

(FILE 'HOME' ENTERED AT 18:27:20 ON 09 MAR 2001)

FILE 'MEDLINE, EMBASE, CAPLUS, BIOSIS' ENTERED AT 18:27:40 ON 09 MAR 2001  
L1 485 S (TALLY, F?)/IN,AU  
L2 855 S (TAO, J?)/IN,AU  
L3 58 S (WENDLER, P?)/IN,AU  
L4 108 S (CONNELLY, G?)/IN,AU  
L5 72 S (GALLANT, C?)/IN,AU  
L6 492 S (GALLANT, D?)/IN,AU  
L7 0 S L1 AND L2 AND L3 AND L4 AND L6  
L8 1977 S L1 OR L2 OR L3 OR L4 OR L6



L9 14 S L8 AND TET  
 L10 6 DUPLICATE REMOVE L9 (8 DUPLICATES REMOVED)  
 L11 15 S L8 AND (METHOD (S) (ISOLATION OR IDENTIFICATION OR SCREENING  
 L12 11 DUPLICATE REMOVE L11 (4 DUPLICATES REMOVED)

=> S L8 AND PRO3

L13 0 L8 AND PRO3

=> S L8 AND PC3844

L14 0 L8 AND PC3844

=> S L8 AND PRORS

L15 0 L8 AND PRORS

=> S L8 AND (INTRODUC? (S) (SUBJECT OR MAMMAL OR ANIMAL OR ORGANISM))

L16 1 L8 AND (INTRODUC? (S) (SUBJECT OR MAMMAL OR ANIMAL OR ORGANISM)  
 )

=> D IBIB AB L16

L16 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1999:451420 CAPLUS

DOCUMENT NUMBER: 131:85158

TITLE: Method for identifying validated target and assay combinations

INVENTOR(S): Tally, Francis P.; Tao, Jianshi; Wendler, Philip A.; Connelly, Gene; Gallant, Paul L.

PATENT ASSIGNEE(S): Cubist Pharmaceuticals, Inc., USA

SOURCE: PCT Int. Appl., 74 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9935494	A1	19990715	WO 1999-US474	19990108
W:				
AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW:				
GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 9922181	A1	19990726	AU 1999-22181	19990108
EP 1046034	A1	20001025	EP 1999-902132	19990108
R:				
AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
NO 2000003515	A	20000907	NO 2000-3515	20000707
PRIORITY APPLN. INFO.:			US 1998-70965	19980109
			US 1998-76638	19980303
			US 1998-81753	19980414
			US 1998-85844	19980518
			US 1998-89828	19980619
			US 1998-94698	19980730
			US 1998-100211	19980914
			US 1998-101718	19980924
			US 1998-107751	19981110

AB The invention comprises methods useful within a laboratory process for identifying compds. and/or designing further compds. with activity to produce a desired phenotype (for example, growth inhibition) in cells whose target cell component is the subject of certain studies to identify such compds. The invention employs constructed cells comprising a regulable gene encoding a biomol. which modulates (inhibits or activates) in vivo the function of a target component of the cell which can be an enzyme for example. The process incorporates methods for identifying biomols. that bind to a chosen target cell component in vitro, methods

for identifying biomols. that also bind to the chosen target and modulate its function intracellularly, causing a phenotypic effect. The intracellular effect of a biomol. can be tested in cell culture, or tested after **introduction** of the constructed cells into a host **mammal** in vivo, and methods for identifying compds. that compete with the biomols. for sites on the target in competitive binding assays. Compds. identified by the series of steps in this process are candidates for

drugs with the desired activity on the cell. Targets for which such compds. can

be identified are validated as being essential to a phenotype of the cell.

REFERENCE COUNT:

2

REFERENCE(S):

(1) Upjohn Co; WO 9117260 A 1991 CAPLUS

(2) Zeneca Ltd; GB 2303209 A 1997 CAPLUS

=> D HIS

(FILE 'HOME' ENTERED AT 18:27:20 ON 09 MAR 2001)

FILE 'MEDLINE, EMBASE, CAPLUS, BIOSIS' ENTERED AT 18:27:40 ON 09 MAR 2001

```
L1      485 S (TALLY, F?)/IN,AU
L2      855 S (TAO, J?)/IN,AU
L3       58 S (WENDLER, P?)/IN,AU
L4      108 S (CONNELLY, G?)/IN,AU
L5       72 S (GALLANT, C?)/IN,AU
L6      492 S (GALLANT, D?)/IN,AU
L7        0 S L1 AND L2 AND L3 AND L4 AND L6
L8     1977 S L1 OR L2 OR L3 OR L4 OR L6
L9       14 S L8 AND TET
L10       6 DUPLICATE REMOVE L9 (8 DUPLICATES REMOVED)
L11      15 S L8 AND (METHOD (S) (ISOLATION OR IDENTIFICATION OR SCREENING
L12     11 DUPLICATE REMOVE L11 (4 DUPLICATES REMOVED)
L13        0 S L8 AND PRO3
L14        0 S L8 AND PC3844
L15        0 S L8 AND PRORS
L16       1 S L8 AND (INTRODUC? (S) (SUBJECT OR MAMMAL OR ANIMAL OR ORGANI
```

=> S ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?) AND (INTRODUC? (S) (SUBJECT OR MAMMAL OR ANIMAL OR ORGANISM))

UNMATCHED LEFT PARENTHESIS '((DRUG'

The number of right parentheses in a query must be equal to the number of left parentheses.

=> S ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?)) AND (INTRODUC? (S) (SUBJECT OR MAMMAL OR ANIMAL OR ORGANISM))

2 FILES SEARCHED...

```
L17     597 ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?)) AND
        (INTRODUC? (S) (SUBJECT OR MAMMAL OR ANIMAL OR ORGANISM))
```

=> S L17 AND TET

L18 0 L17 AND L18

=> S L17 AND INDUC?

L19 126 L17 AND INDUC?

=> S ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?)) AND ((DNA OR VECTOR) (S)INTRODUC? (S) (SUBJECT OR MAMMAL OR ANIMAL OR ORGANISM))

2 FILES SEARCHED...

L20 75 ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?)) AND  
((DNA OR VECTOR) (S) INTRODUC? (S) (SUBJECT OR MAMMAL OR  
ANIMAL  
OR ORGANISM))

=> S ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?)) AND  
((RECOMBINANT (2N) CELL) (S)INTRODUC? (S) (SUBJECT OR MAMMAL OR ANIMAL OR  
ORGANISM))

2 FILES SEARCHED...

L21 0 ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?)) AND  
((RECOMBINANT (2N) CELL) (S) INTRODUC? (S) (SUBJECT OR MAMMAL  
OR ANIMAL OR ORGANISM))

=> S L20 AND (EX (2W) VIVO)

L22 7 L20 AND (EX (2W) VIVO)

=> D TI L22 1-7

L22 ANSWER 1 OF 7 MEDLINE

TI Naked DNA and adenoviral immunizations for immunotherapy of prostate  
cancer: a phase I/II clinical trial.

L22 ANSWER 2 OF 7 MEDLINE

TI Molecular therapy for renal diseases.

L22 ANSWER 3 OF 7 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.

TI Naked DNA and adenoviral immunizations for immunotherapy of prostate  
cancer: A phase I/II clinical trial.

L22 ANSWER 4 OF 7 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.

TI Prostate cancer immunotherapy at the dawn of the new millennium.

L22 ANSWER 5 OF 7 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.

TI Molecular therapy for renal diseases.

L22 ANSWER 6 OF 7 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.

TI The biotechnology of gene therapy.

L22 ANSWER 7 OF 7 BIOSIS COPYRIGHT 2001 BIOSIS

TI Naked DNA and adenoviral immunizations for immunotherapy of prostate  
cancer: A phase I/II clinical trial.

=> D IBIB AB L22 1, 2, 4, 6

L22 ANSWER 1 OF 7 MEDLINE

ACCESSION NUMBER: 2000436262 MEDLINE

DOCUMENT NUMBER: 20355043

TITLE: Naked DNA and adenoviral immunizations for immunotherapy  
of

prostate cancer: a phase I/II clinical trial.

AUTHOR: Mincheff M; Tchakarov S; Zoubak S; Iankov D; Botev C;  
Alankova I; Georgiev G; Petrov S; Iankov H T  
CORPORATE SOURCE: American Foundation for Biological Research, Rockville, MD  
20852, USA.. mincheffm@netscape.net  
SOURCE: EUROPEAN UROLOGY, (2000 Aug) 38 (2) 208-17.  
Journal code: ENM. ISSN: 0302-2838.  
PUB. COUNTRY: Switzerland  
(CLINICAL TRIAL)  
(CLINICAL TRIAL, PHASE I)  
(CLINICAL TRIAL, PHASE II)  
Journal; Article; (JOURNAL ARTICLE)  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 200011  
ENTRY WEEK: 20001104

AB **INTRODUCTION AND OBJECTIVES:** Animal studies have indicated that the use of syngeneic dendritic cells that have been transfected **ex vivo** with **DNA** for tumor-specific antigen results in tumor regression and decreased number of metastases. Additional studies have also suggested the possibility to modulate the dendritic cells in vivo either by 'naked' **DNA** immunization or by injecting replication-deficient viral vectors that carry the tumor-specific **DNA**. Using the prostate-specific membrane antigen (PSMA) as a **target** molecule, we have initiated a clinical trial for immunotherapy of prostate cancer. The primary objective of the study was to **determine** the safety of the PSMA vaccine after repeated intradermal injections. **METHODS:** We have included the extracellular human PSMA **DNA** as well as the human CD86 **DNA** into separate expression vectors (PSMA and CD86 plasmids), and into a combined PSMA/CD86 plasmid. In addition, the expression cassette from the PSMA plasmid was inserted into a replication deficient adenoviral expression **vector**. Twenty-six patients with prostate cancer were entered into a phase I/II toxicity-dose escalation study, which was initiated in spring 1998. Immunizations were performed intradermally at weekly intervals. Doses of **DNA** between 100 and 800  $\mu$ g and of recombinant virus at  $5 \times 10^8$  PFUs per application were used. **RESULTS AND CONCLUSION:** No immediate or long-term side effects following immunizations have been recorded. All patients who received initial inoculation with the viral **vector** followed by PSMA plasmid boosts showed signs of immunization as evidenced by the development of a delayed-type hypersensitivity reaction after the PSMA plasmid injection. In contrast, of the patients who received a PSMA plasmid and CD86 plasmid, only 50% showed signs of successful immunization. Of the patients who received PSMA plasmid and soluble GM-CSF, 67% were immunized. However, all patients who received the PSMA/CD86 plasmid and sGM-CSF became immunized. The patients who did not immunize during the first round were later successfully immunized after a boost with the viral **vector**. The heterogeneity of the medical status and the presence in many patients of concomitant hormone therapy does not permit unequivocal interpretation of the data with respect to the effectiveness of the therapy. However, several responders, as evidenced by a change in the local disease, distant metastases, and PSA levels, can be **identified**. A phase II clinical study to evaluate the effectiveness of the therapy is currently underway.

L22 ANSWER 2 OF 7 MEDLINE  
ACCESSION NUMBER: 96438576 MEDLINE  
DOCUMENT NUMBER: 96438576  
TITLE: Molecular therapy for renal diseases.  
AUTHOR: Lipkowitz M S; Klotman M E; Bruggeman L A; Nicklin P; Hanss

CORPORATE SOURCE: B; Cappaport J; Klotman P E  
Department of Medicine, Mount Sinai School of Medicine,  
New

York, NY 10029, USA.  
SOURCE: AMERICAN JOURNAL OF KIDNEY DISEASES, (1996 Oct) 28 (4)  
475-92. Ref: 169  
Journal code: 3H5. ISSN: 0272-6386.

PUB. COUNTRY: United States  
Journal; Article; (JOURNAL ARTICLE)  
General Review; (REVIEW)  
(REVIEW, TUTORIAL)

LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199701  
ENTRY WEEK: 19970104

AB The **introduction** of molecular therapy through the delivery of nucleic acids either as oligonucleotides or genetic constructs holds enormous promise for the treatment of renal disease. Significant barriers remain, however, before successful organ-specific molecular therapy can be applied to the kidney. These include the development of methods to **target** the kidney selectively, the definition of vectors that transduce renal tissue, the **identification** of appropriate molecular targets, the development of constructs that are regulated and expressed for long periods of time, the demonstration of efficacy in vivo, and the demonstration of safety in humans. As the genetic and pathophysiologic basis of renal disease is clarified, obvious targets for therapy will be defined, for example, polycystin in polycystic kidney disease, human immunodeficiency virus (HIV) type 1 in HIV-associated nephropathy, alpha-galactosidase A in Fabry's disease, insulin in diabetic nephropathy, and the "minor" collagen IV chains in Alport's syndrome. In addition, several potential mediators of progressive renal disease may be amenable to molecular therapeutic strategies, such as interleukin-6, basic fibroblast growth factor (bFGF), platelet-derived growth factor (PDGF), and transforming growth factor-beta (TGF-beta). To test the in vivo efficacy of molecular therapy, appropriate **animal** models for these disease states must be developed, an area that has received too little attention. For the successful delivery of genetic constructs to the kidney, both viral and nonviral **vector** systems will be required. The kidney has a major advantage over other solid organs since it is accessible by many routes, including intrarenal artery infusion, retrograde delivery through the uroexcretory pathways, and **ex vivo** during transplantation. To further restrict expression to the kidney, tropic vectors and tissue-specific promoters also must be developed. For the purpose of inhibition of endogenous or exogenous genes, current therapeutic modalities include the delivery of antisense oligodeoxynucleotides or ribozymes. For these approaches to succeed, we must gain a much better understanding of the nature of their transport into the kidney, requirements for specificity, and in vivo mechanisms of action. The danger of a rush to clinical application is that superficial approaches to these issues will likely fail and enthusiasm will be lost for an area that should be one of the most exciting developments in therapeutics in the next decade.

L22 ANSWER 4 OF 7 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.  
ACCESSION NUMBER: 2000197685 EMBASE  
TITLE: Prostate cancer immunotherapy at the dawn of the new millennium.  
AUTHOR: Salgaller M.L.  
CORPORATE SOURCE: M.L. Salgaller, Northwest Biotherapeutics, Inc., 2203 Airport Way South, Seattle, WA 98134, United States.

ml.nwbio.com  
SOURCE: Expert Opinion on Investigational Drugs, (2000) 9/6  
(1217-1229).  
Refs: 108  
ISSN: 1354-3784 CODEN: EOIDER  
COUNTRY: United Kingdom  
DOCUMENT TYPE: Journal; General Review  
FILE SEGMENT: 003 Endocrinology  
006 Internal Medicine  
016 Cancer  
026 Immunology, Serology and Transplantation  
030 Pharmacology  
037 Drug Literature Index  
038 Adverse Reactions Titles  
LANGUAGE: English  
SUMMARY LANGUAGE: English

AB Standard treatments for adenocarcinoma of the prostate, such as surgery, hormones, radiation and chemotherapy, often achieve a clinical response, but this is usually short-lived. Prostate cancer frequently recurs and second-line therapies have a poor response rate. Many clinicians seem comfortable in limiting their philosophy of treating advanced recurrent disease merely to new regimens of failed therapies, such as combination chemotherapy. However, other medical researchers have chosen to pursue novel approaches, including immunotherapy, several of which are summarised in this review. Although ranging widely in antigen specificity, all attempt to exploit the body's natural antitumour immunity. Furthermore, all aim to stimulate immunity above a threshold level necessary for tumour regression or to induce stability in the face of progression. The goal of in vivo or **ex vivo** gene therapy is the modification of gene expression within an antigen-presented cell by the **introduction** of a **vector**, **DNA**, or RNA. Within that field, much progress has been made and is ongoing currently concerning gene delivery systems, **target identification** and characterisation. Comparatively, monoclonal antibodies are an established type of cancer immunotherapy. However, the more recent development of humanised or fully human antibodies, as well as novel moieties they can be coupled to, renews their prospects for clinical impact. Lastly, various cell-based therapies are the focus of several recent clinical studies demonstrating tumour regression or stabilisation. Immune cells, for example, T-lymphocytes and dendritic cells, have already demonstrated treatment benefit, as well as the ability to maintain an excellent quality of life for participants. Overall, there is a multitude of approaches being considered for the treatment of prostate cancer. The following review concentrates on those approaches that are currently in human or **animal** studies and have a specific emphasis on prostate cancer.

L22 ANSWER 6 OF 7 EMBASE COPYRIGHT 2001 ELSEVIER SCI. B.V.  
ACCESSION NUMBER: 96251000 EMBASE  
DOCUMENT NUMBER: 1996251000  
TITLE: The biotechnology of gene therapy.  
AUTHOR: Pappas M.G.  
CORPORATE SOURCE: Advanced Instruments, Inc., Two Technology Way, Norwood, MA  
02062, United States  
SOURCE: Drug Development and Industrial Pharmacy, (1996) 22/8  
(791-803).  
ISSN: 0363-9045 CODEN: DDIPD8  
COUNTRY: United States  
DOCUMENT TYPE: Journal; General Review  
FILE SEGMENT: 004 Microbiology  
005 General Pathology and Pathological Anatomy  
016 Cancer  
022 Human Genetics

LANGUAGE: English

SUMMARY LANGUAGE: English

AB The prospect for correcting highly morbid or fatal inherited diseases, or ameliorating cancer and acquired, deadly infectious diseases such as AIDS using gene therapy is very exciting. Numerous recent advances in molecular

biology make it possible, not only to **identify** and locate genes associated with human inherited disorders and cancers, but to potentially correct these disorders with functional genes. These advances include

more rapid gene **identification**, isolation and sequencing techniques, a better understanding of the functions and relationships between genes and their products in vivo, the development and study of human and model **organism** genomes, elucidation of genetic disease pathology using **animal** genetic disease models, advanced computer amino acid and nucleotide sequencing software and data bases, and the development and

use of novel chemical, physical, and viral **vector** gene delivery methods. Functional genes are **introduced** using two approaches, **ex vivo** and in vivo gene therapy. In **ex vivo** therapy, autologous cells are removed from the patient, genetically altered by inserting the functional gene, characterized, and then returned to the patient; in in vivo therapy, functional genes are packaged for delivery directly into the patient, where cellular uptake

and gene expression occurs. Scores of clinical trials have been federally approved to treat patients with a variety of inherited disorders, cancers,

and acquired diseases using these two approaches. Roadblocks to long-lasting gene therapy include understanding more completely the biological functions of somatic cells or organs targeted for gene

therapy, targeting appropriate host cells and achieving high gene delivery rates

in these cells, regulating and sustaining gene expression through optimal **DNA** insertion into chromosomes such that other cellular functions are not adversely affected, and the prevention of **vector**-induced diseases or cancers. Ethical considerations regarding proper use of somatic gene therapy and the potential for germline gene therapy must

also be seriously considered. The prospect of permanent correction of highly morbid or fatal maladies using gene therapy could prove to be one of the great advances in public health and could revolutionize the **identification** and gene-**drug** treatment of a broad spectrum of inherited and acquired human diseases.

=&gt; D HIS

(FILE 'HOME' ENTERED AT 18:27:20 ON 09 MAR 2001)

FILE 'MEDLINE, EMBASE, CAPLUS, BIOSIS' ENTERED AT 18:27:40 ON 09 MAR 2001

L1 485 S (TALLY, F?)/IN,AU  
 L2 855 S (TAO, J?)/IN,AU  
 L3 58 S (WENDLER, P?)/IN,AU  
 L4 108 S (CONNELLY, G?)/IN,AU  
 L5 72 S (GALLANT, C?)/IN,AU  
 L6 492 S (GALLANT, D?)/IN,AU  
 L7 0 S L1 AND L2 AND L3 AND L4 AND L6  
 L8 1977 S L1 OR L2 OR L3 OR L4 OR L6  
 L9 14 S L8 AND TET  
 L10 6 DUPLICATE REMOVE L9 (8 DUPLICATES REMOVED)  
 L11 15 S L8 AND (METHOD (S) (ISOLATION OR IDENTIFICATION OR SCREENING  
 L12 11 DUPLICATE REMOVE L11 (4 DUPLICATES REMOVED)

L13 0 S L8 AND PRO3  
L14 0 S L8 AND PC3844  
L15 0 S L8 AND PRORS  
L16 1 S L8 AND (INTRODUC? (S) (SUBJECT OR MAMMAL OR ANIMAL OR ORGANI  
L17 597 S ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?))  
AND  
L18 0 S L17 AND TET  
L19 126 S L17 AND INDUC?  
L20 75 S ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?))  
AND  
L21 0 S ((DRUG OR TARGET) (S) (IDENTIF? OR SCREEN? OR DETERMIN?))  
AND  
L22 7 S L20 AND (EX (2W) VIVO)



EAST: DERWENT EPO JPO USPAT 3/11/01 09/227,687

L1 12 tally-f\$.in.  
L2 119 tao-j\$.in.  
L3 32 wendler-p\$.in.  
L4 11 connelly-g\$.in.  
L5 78 gallant-d\$.in.  
L6 154 shen-x\$.in.  
L7 1820 zhang-j\$.in.  
L8 2211 1 or 2 or 3 or 4 or 5 or 6 or 7  
L9 7 8 and aureus  
L10 133 methionyl\$ and aureus  
L11 9 (methionyl\$ with synthetase) and aureus  
L12 1069 pathogen and (method with (screening or identifying or isolating or determining) with (compound or target or inhibitor))  
L13 313 12 and aureus  
L14 4 13 and (methionyl\$ with synthetase)  
L15 327 pathogen and (method with (screening or identifying or isolating or determining) with (compound or target) with inhibit\$)  
L16 0 15 and (in adj2 vivo)  
L17 5 15 and (test adj3 animal)  
L18 10 pathogen and (method with (screening or identifying or isolating or determining) with (compound or target) with inhibit\$) with animal  
L19 10 pathogen and (method with (screening or identifying or isolating or determining) with (compound or target) with inhibit\$ with animal)  
L20 10 18 or 19  
L21 109 pathogen and (method with (screening or identifying or isolating or determining) with (compound or target) with inhibit\$) and aureus  
L22 159 pathogen and ( (screening or identifying or isolating or determining) with (compound or target) with inhibit\$) and aureus  
L23 16 pathogen and ( (screening or identifying or isolating or determining) with (compound or target) with inhibit\$ with animal) and aureus  
L24 309 pathogen and ( (screening or identifying or isolating or determining) with (compound or target) with inhibit\$) and (animal with (introduc\$ or infect\$))  
L25 235 pathogen and ( (screening or identifying or isolating or determining) with (compound or target) with inhibit\$) and (animal with (introduc\$ or infect\$) with (cell or pathogen))  
L26 84 pathogen and ( (screening or identifying or isolating or determining) with (compound or target) with inhibit\$) and (animal with (introduc\$ or infect\$) with (cell or pathogen)) and aureus  
L27 0 pathogen and ( (screening or identifying or isolating or determining) with (compound or target) with inhibit\$ with animal with (introduc\$ or infect\$) with (cell or pathogen)) and aureus